



## NPHS1 gene

NPHS1, nephrin

### Normal Function

The *NPHS1* gene provides instructions for making a protein called nephrin. Nephrin is primarily found in the kidneys, which are organs that filter waste products from the blood and remove them in urine. Specifically, nephrin is found in cells called podocytes, which are located in specialized kidney structures called glomeruli. Nephrin is located at the cell surface in the area between two podocytes called the slit diaphragm. The slit diaphragm is known as a filtration barrier because it captures proteins in blood so that they remain in the body while allowing other molecules like sugars and salts to be excreted in urine. Nephrin proteins on one cell interact with nephrin proteins on adjacent podocytes, forming a zipper-like structure. This structure allows the passage of small molecules through the slit diaphragm while preventing larger molecules like proteins from passing through. Nephrin proteins are essential for forming the slit diaphragm, anchoring the slit diaphragm to podocytes, and filtering blood.

Nephrin is also involved in cell signaling. It relays signals from outside the cell to inside the cell. Additionally, nephrin proteins on the surface of adjacent cells send and receive signals, allowing podocytes to communicate with one another.

### Health Conditions Related to Genetic Changes

#### congenital nephrotic syndrome

At least 250 mutations in the *NPHS1* gene have been found to cause congenital nephrotic syndrome. This condition is a kidney disorder that begins in infancy and typically leads to irreversible kidney failure (end-stage renal disease) by early childhood.

*NPHS1* gene mutations account for all cases of congenital nephrotic syndrome of the Finnish type. This form of the condition is found in people of Finnish ancestry. Two specific mutations, both of which result in an abnormally short, nonfunctional nephrin protein, account for nearly all cases. The first mutation, known as Finn<sub>major</sub>, is written as L41fsX90 and is responsible for 78 percent of cases. The second mutation, known as Finn<sub>minor</sub>, is written as R1109X and is responsible for 16 percent of cases.


*NPHS1* gene mutations can also cause congenital nephrotic syndrome in non-Finnish individuals. Most of these mutations result in an abnormal nephrin protein that is trapped inside the cell and cannot get to the podocyte cell surface. A shortage of functional nephrin at the podocyte cell surface impairs the formation of normal slit

diaphragms. Without a functional slit diaphragm, molecules pass through the kidneys abnormally and are excreted in urine. The filtering ability of the kidneys worsens from birth, eventually leading to end-stage renal disease.

### Chromosomal Location

Cytogenetic Location: 19q13.12, which is the long (q) arm of chromosome 19 at position 13.12

Molecular Location: base pairs 35,825,372 to 35,851,993 on chromosome 19 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

### Other Names for This Gene

- CNF
- nephrin
- nephrin precursor
- nephrosis 1, congenital, Finnish type (nephrin)
- NPHN
- NPHS1 nephrin
- renal glomerulus-specific cell adhesion receptor

### Additional Information & Resources

#### Educational Resources

- National Institute of Diabetes and Digestive and Kidney Diseases: The Kidneys and How They Work  
<https://www.niddk.nih.gov/health-information/kidney-disease/kidneys-how-they-work>

### Scientific Articles on PubMed

- PubMed  
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28NPHS1%5BTIAB%5D%29+OR+%28nephrin%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1080+days%22%5Bdp%5D>

### OMIM

- NEPHRIN  
<http://omim.org/entry/602716>

### Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology  
[http://atlasgeneticsoncology.org/Genes/GC\\_NPHS1.html](http://atlasgeneticsoncology.org/Genes/GC_NPHS1.html)
- ClinVar  
<https://www.ncbi.nlm.nih.gov/clinvar?term=NPHS1%5Bgene%5D>
- HGNC Gene Family: C2-set domain containing  
<http://www.genenames.org/cgi-bin/genefamilies/set/592>
- HGNC Gene Family: Fibronectin type III domain containing  
<http://www.genenames.org/cgi-bin/genefamilies/set/555>
- HGNC Gene Family: Immunoglobulin like domain containing  
<http://www.genenames.org/cgi-bin/genefamilies/set/594>
- HGNC Gene Family: V-set domain containing  
<http://www.genenames.org/cgi-bin/genefamilies/set/590>
- HGNC Gene Symbol Report  
[http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?q=data/hgnc\\_data.php&hgnc\\_id=7908](http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=7908)
- NCBI Gene  
<https://www.ncbi.nlm.nih.gov/gene/4868>
- UniProt  
<http://www.uniprot.org/uniprot/O60500>

## Sources for This Summary

- Fogo AB, Lusco MA, Najafian B, Alpers CE. AJKD Atlas of Renal Pathology: Congenital Nephrotic Syndrome of Finnish Type. Am J Kidney Dis. 2015 Sep;66(3):e11-2. doi: 10.1053/j.ajkd.2015.07.008.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/26300201>
- Grahammer F, Schell C, Huber TB. The podocyte slit diaphragm--from a thin grey line to a complex signalling hub. Nat Rev Nephrol. 2013 Oct;9(10):587-98. doi: 10.1038/nrneph.2013.169. Epub 2013 Sep 3. Review.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/23999399>
- OMIM: NEPHRIN  
<http://omim.org/entry/602716>
- Ovunc B, Ashraf S, Vega-Warner V, Bockenhauer D, Elshakhs NA, Joseph M, Hildebrandt F; Gesellschaft für Pädiatrische Nephrologie (GPN) Study Group. Mutation analysis of NPHS1 in a worldwide cohort of congenital nephrotic syndrome patients. Nephron Clin Pract. 2012;120(3):c139-46. doi: 10.1159/000337379. Epub 2012 May 11.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/22584503>
- Schoeb DS, Chernin G, Heeringa SF, Matejas V, Held S, Vega-Warner V, Bockenhauer D, Vlangos CN, Moorani KN, Neuhaus TJ, Kari JA, MacDonald J, Saisawat P, Ashraf S, Ovunc B, Zenker M, Hildebrandt F; Gessellschaft für Paediatric Nephrologie (GPN) Study Group. Nineteen novel NPHS1 mutations in a worldwide cohort of patients with congenital nephrotic syndrome (CNS). Nephrol Dial Transplant. 2010 Sep;25(9):2970-6. doi: 10.1093/ndt/gfq088. Epub 2010 Feb 18.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/20172850>  
*Free article on PubMed Central:* <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2948833/>
- Wang JJ, Mao JH. The etiology of congenital nephrotic syndrome: current status and challenges. World J Pediatr. 2016 May;12(2):149-58. doi: 10.1007/s12519-016-0009-y. Epub 2016 Mar 9.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/26961288>

---

Reprinted from Genetics Home Reference:  
<https://ghr.nlm.nih.gov/gene/NPHS1>

Reviewed: July 2016

Published: March 21, 2017

Lister Hill National Center for Biomedical Communications  
U.S. National Library of Medicine  
National Institutes of Health  
Department of Health & Human Services